

Supplementary file for: Kilias E.S., Peeken I. & Metfies K. 2014. Insight into protist diversity in Arctic sea ice and melt-pond aggregate obtained by pyrosequencing. *Polar Research 33*. Correspondence: E.S. Kilias, Alfred Wegener Institute for Polar and Marine Research, Am Handelshafen 12, DE-27570 Bremerhaven, Germany. E-mail: estelle.kilias@awi.de.



Supplementary Fig. S1. Aggregates accumulating at the bottom of the melt pond. Insert shows an enlarged image of the sampled aggregates.



Supplementary Fig. S2. Phylogenetic tree of the abundant biosphere, computed on the basis of the V4 region for the consensus (square), original (circle) and BLAST search sequences (triangle). The optimal nucleotide substitution model was identified by MODELGENERATOR, while the phylogenetic tree was constructed in MEGA (GTR+I+G).

Supplementary Table S1. Taxonomic annotation (BLAST search; National Center for Biotechnology Information) of consensus and original sequences, showing identical identifications with similar qualities for both sequence types. Phylotype ID of the abundant biosphere ( $\geq 1\%$ ) was used from a detailed ARB-based annotation. Consensus sequences were obtained from a sequence clustering on a similarity threshold of 97% (DNAStar). Original sequences were abundant contributors to the sequence clusters (operational taxonomic unit).

| Phylotype ID                    | consensus sequence |          |                  | original sequence |          |                  |
|---------------------------------|--------------------|----------|------------------|-------------------|----------|------------------|
|                                 | Query<br>cover     | Identity | Accession number | Query<br>cover    | Identity | Accession number |
| Unc. <sup>a</sup> Gymnodiniales | 98                 | 99       | KC488417.1       | 100               | 99       | KC488417.1       |
| Peridinium sp.                  | 100                | 99       | KF446624.1       | 100               | 99       | KF446624.1       |
| Strombidium basimorphum         | 99                 | 100      | AJ431358.1       | 100               | 99       | AJ431358.1       |
| Carteria sp.                    | 98                 | 100      | JN934685.1       | 100               | 99       | JN934685.1       |
| Chlamydomonadaceae 1            | 98                 | 99       | EF432529.1       | 99                | 99       | EF432529.1       |
| Chlamydomonadaceae 2            | 99                 | 98       | EF432529.1       | 99                | 98       | EF432529.1       |
| Chlamydomonadaceae 3            | 98                 | 99       | FN690710.1       | 100               | 99       | FN690710.1       |
| Chlamydomonadaceae 4            | 99                 | 98       | FN690710.1       | 99                | 97       | FN690710.1       |
| Dinobryon faciluiferum 1        | 97                 | 99       | EU545761.1       | 98                | 99       | EU545761.1       |
| Dinobryon faciluiferum 2        | 100                | 98       | JF730832.1       | 100               | 97       | JF730832.1       |
| Ochromonas tuberulata           | 98                 | 96       | AB4749151.1      | 99                | 96       | AB4749151.1      |
| Ochromonas sp. 1                | 98                 | 100      | EU247834.1       | 100               | 99       | EU247834.1       |
| Ochromonas sp. 2                | 99                 | 99       | EF165133.1       | 100               | 99       | EF165133.1       |
| Ochromonas sp. 3                | 100                | 98       | EU076745.1       | 99                | 92       | EU076745.1       |
| Pelagomonas sp.                 | 100                | 100      | EU247837.1       | 99                | 99       | EU247837.1       |

<sup>&</sup>lt;sup>a</sup> Uncultured.